## RESEARCH AND EVALUATION OF THE FUNCTIONAL CAPABILITIES AND PHYSICAL DEVELOPMENT OF QUALIFIED BOXERS UNDER THE INFLUENCE OF THE TRAINING PROCESS

Docent **Vladimir Serebryakov Vladimirovich**Uzbek State University of Physical Culture and Sports

Senior teacher **Khushvakov Nurxon Yusupovich** Uzbek State University of Physical Culture and Sports

Teacher **Asatillaev Jokhongir Ne'matjon ugli**Uzbek State University of Physical Culture and Sports

**Annotation.** The article presents an assessment of the physical development and functional capabilities of qualified boxers, which was carried out under the influence of training loads. Similarities and differences in physique indicators are also determined, which is directly related to the specifics of the training process of the compared specializations. It is concluded that in boxing, the loads are more intense in nature, since this sport involves the maximum return in short periods of time with a relatively slow recovery of energy costs.

**Key words**: physique, cardiorespiratory system, bicycle egometric test, anthropometry, aerobic system, physical activity.

Relevance. When developing model characteristics of athletes for the purpose of selection, in addition to a number of physique indicators, data on the composition of body weight, functional reserves of the cardiorespiratory system, the indicators of which are an indicator of sports performance, are of particular interest. The fundamental position of sports selection is the idea of population reality of individuals with extremely high severity of morphological and functional qualities, the importance of which is undeniable for achieving success in sports (V.B. Schwartz, 1991, B.A. Nikityuk, 1996, Rogozkin V.A., Nazarov I.B. et al., 2006, Safarova D.D. et al., 2009, 2018). In this regard, the morpho-functional indicators of boxers can be used not only to identify the state of fitness of athletes, but it is possible to clearly plan the direction of the training process, as well as correctly and objectively conduct sports selection. In order to obtain additional information about the effectiveness of the training process not only pedagogical, but also physiological indicators of the sides of the preparedness of athletes were recorded in a number of tests. In the work, in addition to studying the

April, 2023

morphological features of athletes, the task was to find objective criteria for determining physical fitness based on the functional state of athletes at different stages of the training process, to achieve optimal readiness (peak of sports form) for competitions and assess the state of rehabilitation after sports competitions. These problems were and remain topical issues at the present time. One of the most important indicators of the reserve capacity of the body is the state of the cardio-respiratory system,

Purpose of the study: Assessment of physical development and functional readiness of qualified boxers on the basis of anthropometry signs and indicators of the cardiorespiratory system.

**Research objectives:** To assess some indicators of the cardiorespiratory system, a survey of athletes studying at the Uzbek State University of Physical Culture was conducted. The study involved qualified boxers with the title of CMS (8) and 1 category (2). The total number of examined athletes was 10 young men aged 18-20 years. To conduct research in order to determine the physical development and characteristics of the cardiorespiratory system, the following anthropometric indicators were measured in the subjects: height, weight, chest circumference at rest (RGC). In addition, the vital capacity of the lungs (VC), arterial pressure - systolic (SD) and diastolic (DD) were determined. Heart rate (HR) was determined by pulse. A hypoxic test was also carried out (the time of maximum delay inspiratory breath measured in seconds). As a load test, we chose a 9-minute bicycle ergometer test. All data obtained were statistically processed and presented in the form of tables, which also indicate the significance of each indicator.

## Research results and discussion:

To assess physical development, data are presented on intragroup variability in the variability of total body sizes in acyclic sports - boxers, fencers (Table 1).

The greatest body length is observed in fencers - 177.14  $\pm$  7.19 cm, in boxers - $173 \pm 5.74$  cm.  $13\pm 4.43$  kg.

The reverse trend is typical for the average values of chest girth for boxers -93.55±6.54 cm, for fencers - 90.67±4.35 cm. The values of standard deviations range from 4.43 to 8.93. When assessing the total body dimensions of athletes, it should be noted that boxers with a smaller body length and body weight have the largest chest and hip measurements.

Among the partial signs, representatives of the 2 specializations revealed significant differences in the length of the thigh, lower leg, neck, abdomen, and

April, 2023

shoulder girth. According to these signs, as well as in a number of other anthropometric indicators, the samples of boxers and fencers seemed to be more similar.

Table 1
Indicators of the total body size of athletes specializing in acyclic sports

№	signs	Options	Boxing n=24	Fencing n=20
1	body length	S±m Dispersion Wed sq. deviation	173.61±0.77 32.95 5.74	177.14±1.61 51.69 7.19
2	Body mass	S±m Dispersion Wed sq. deviation	64.13±0.91 45.85 6.77	65.64±2.00 79.60 8.93
3	Chest girth- noah cells	S±m Dispersion Wed sq. deviated knowledge	93.55 42.79 6.54	90.60±0.97 18.89 4.35

Table 2 presents all indicators of the cardiorespiratory system before and after exercise.

 $\begin{tabular}{l} \it Table~2\\ \it Indicators~of~the~cardiorespiratory~system~before~physical~activity~and~after~physical~activity~M\pm m \end{tabular}$ 

№	Options	Indicators up to physical activity (n=10 boxers)	Indicators after physical activity (n=10 boxers)
1	height	172.3±3.1	172.3±3.1
2	weight	71.6±4.6	71.6±4.6
3	chest girth	90±3.5	95±3.4
4	VC, ml	4.4±1.2	4.2±1.2
5	Breath holding on	70.1±3.5	66.4±3.9
	inspiration, s		
6	Heart rate in 1 min	69.4±1.3	85.1±3.1
7	Arterial pressure, max	120±1.2	120.5±3.8
8	Arterial pressure, min	72.5±3.4	68.12.1

Table 2 presents data on indicators of VC and breath holding on inspiration for 1 minute, which made it possible to evaluate the aerobic capabilities of boxers in a comparative aspect.

**Conclusions:** The data obtained allowed us to draw certain conclusions and formulate the following research results:

- a) in qualified boxers, the starting capabilities of the cardiorespiratory system are quite high and stable and are within the normal range;
- b) after intense physical activity, cardiorespiratory indicators also increase within the normal range and return to the original data in a relatively short period of time;
- c) the degree and index of recovery in qualified boxers are quite high, which indicates endurance and a high level of fitness.

Similarities and differences in physique indicators have been established, which directly related is the specifics of the process the to training compared specializations. In addition, it can be assumed that in boxing, the loads are more intense in nature, since this sport involves the maximum return in short periods of time with relatively slow recovery energy costs.

## **LITERATURE**

1. Абрамова Т.Ф., Никитина Т.М., Кочеткова Н.И., Арьков В.В. и др.

Особенности телосложения функционального оценке состояния спортсменов

/ Межд. Конф. Проблемы современной морфологии человека М., 2008 стр. 127.

- 2. Дорохов Р.Н. Основа и перспектива возрастного соматотипирования. // "Теория и практика физической культуры", Москва, 2000, №9, с. 10.
- 3. Ковальчук Г.И. Прогнозирование двигательных способностей бегунов на короткие дистанции. // "Теория и практика физической культуры", Москва, 2003, №9, c. 31-55.
- 4. Никитюк Б.А. Адаптация, конституция, моторика //Теория и практика физической культуры №1, 1989, стр. 40 – 42
- 5. Павлова О.И. Комплексная система подготовки квалифицированных легкоатлетов на этапах становления спортивного мастерства, Москва, 2004, -119 стр.

April, 2023

- 6. Панасюк Т.В., Тамбовцева Р.В. Конституциональные особенности физической работоспособности подростков. В сб. VII Международный науч. конгресс "Современный Олимпийский спорт и спорт для всех", том 2, Москва 2003, c. 135-136.
  - 7. Сафарова Д.Д. Спорт морфологияси (дарслик) Ташкент, 2015, 204 с.
- 8. J.E. Lindsay Carter, Barbara Howeyman Heath Somatotyping development and applications./Department of physical Education San-Diego California, USA, Department of Antropology, Pennsylvania, USA. (Cambridge university press, 1989). -320 c.
- 9. Cova E. P. Human variation in motor abilities and its genetic analisis Cherles Univ. 1981, Pragae, 178 p
- 10. Олейник, Н.А., Чибисов, В.И., Рейдерман, Ю.И., Скрипченко, И.Т., Чередник, Е.А. (2012), "Разработка методики определения текущего состояния сердечно-сосудистой и дыхательной систем у спортсменов", Слобожанський науково-спортивний вісник, № 5-2
  - (33), C. 92-96.
- 11. Сайёра Саъдуллаевна Шукурова, Мавлюда Джалаловна Пулатова, Владимир Владимирович Серебряков; "Изменения показателей макроэлементов в крови у футболистов после физической нагрузки" Academic research in educational sciences 2

(Special Issue 1), 278-286, 2021

- 12. Серебряков.В.В, Хушваков.Н.Ю "Спортивная Физиология" ўкув кўлланма 2023, 152c
- 13. Пулатова Мавлуда Джалоловна "Спорт физиологияси" ўкув кўлланма 2023, 85c
- 14. Сейдалиева Лейла Джанибековна, Серебряков Владимир Владимирович, Мусаева Умида Ахмедовна "forming a healthy lifestyle in physical culture lessons" 2019
- 15. Романов, Ю.Н., Аминов, А.С., Романова, Л.А. (2016), "Современные методики мониторинга функционального состояния кардиопульмональной системы. обшей и специальной работоспособности боксеров квалификации", Человек. Спорт. Медицина, Т. 1, № 1, С. 34-41.